


RESTRICTED USE PESTICIDE

Due to acute oral and inhalation toxicity.

For retail sale and application only by certified applicators or personnel under their direct supervision.

EPA SLN NO.

MT-860007



FOR DISTRIBUTION
AND USE ONLY IN
MONTANA

Furadan® 4F

Insecticide-Nematicide

For Agricultural or Commercial Use Only

EPA REG. No. 279-2876 ZA

EMERGENCY CALLS: 800-331-3148

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN
A MANNER INCONSISTENT WITH ITS LABELING.

CROP	PEST	RATE OF APPLICATION
Wheat	Grasshoppers	0.25 to 0.5 fluid ounces per 1,000 linear feet of row

DIRECTIONS FOR USE:

For planting time application for early season seedling grasshopper control, use Furadan 4F at a rate of 0.25 to 0.5 fluid ounces per 1,000 linear feet of row with a minimum row spacing of 6 inches. Apply into the seed furrow using a microtube or suitable application method to concentrate the Furadan 4F solution around the seed. The use of the higher Furadan 4F rates is recommended where moderate to heavy grasshopper populations exist or where an extended protection period is desired.

Do not use this product through any type of irrigation system.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

PERSONAL PROTECTIVE EQUIPMENT AND REENTRY INTERVALS ON THIS 24C OR ON THE BASIC LABEL MUST BE FOLLOWED.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

24(c) Registrant:

Code 279/2837 3/94

FMC Corporation
Agricultural Chemical Group
1735 Market Street
Philadelphia, PA 19103

MT-860007

FMC®

Furadan and **FMC** - FMC Trademarks
©1991 FMC Corporation